

Inga fagifolia* (L.) Willd.*Guamá**SO-ITF-SM-72
April 1994**Leguminosae (Mimosoideae)****Legume family****John K. Francis**

Inga fagifolia (L.) Willd., commonly known as guamá (Spanish), sweet-pea and Spanish oak (English), and pois doux (French), is a medium-sized (fig. 1) evergreen tree native to the Greater and Lesser Antilles. It is common in moist and wet forests, especially secondary forests. The tree is still used to some extent as a coffee shade tree and is sawed for lumber.

HABITAT**Native Range**

The native range of guamá (fig. 2) includes the Greater Antilles, except Cuba, where it is naturalized (5); the Virgin Islands; Antigua; Saba; St. Kitts; Montserrat; Guadeloupe; Marie Galante; Dominica; Martinique; St. Vincent; Grenada; Barbados in the Lesser Antilles; and Trinidad (13, 16). The distribution usually ascribed to guamá in Venezuela (24) may actually be that of other species of *Inga*, as is the distribution of what was once thought to be guamá in Mexico and Central America (13).

Climate

Guamá is adapted to moist and wet forests, with mean annual rainfall from 1400 to 3500 mm. The species also grows in drier areas on alluvial soils along perennial or intermittent streams. Dry seasons of up to 3 months are a feature of the climate of most of the islands where guamá grows. Mean annual temperatures within the native range of guamá range from about 23 to 26 °C, depending mainly on altitude. Temperatures rarely exceed 32 °C or drop below 15 °C. Frosts are unknown in the native range of guamá.

Soils and Topography

Guamá is able to grow on a wide variety of soils and sites, including soils with textures from sands to clays (20). The species grows on soils originating from many kinds of parent material. Soil pH at least as low as 5.0 and moderately low levels of exchangeable cations seem to be adequate for guamá. Soils can be poorly drained, but not swampy. Excessively drained soils due to sand or skeletal rock or shallow soils in areas of minimum rainfall will not support the species. Guamá grows on both steep and level topography from a few meters above sea level to over 1,000 m in elevation.

Associated Forest Cover

In a subtropical wet forest stand dominated by *Dacryodes excelsa* Vah., *Euterpe globosa* Gaertn., *Cecropia schreberiana* Miq., *Micropholis garcinifolia* Pierre, *Slonea berteriana* Choisy, *Cyrilla racemiflora* L., and *Magnolia splendens* Urban, guamá contributed only 72 of a total of 3,100 stems (25). In another subtropical wet forest dominated by *C. racemiflora* L., *Micropholis garcinifolia*, and *Magnolia splendens*, guamá contributed 28 of 3,400 stems (25). Guamá, *Inga vera* Willd., *Guarea guidonia* (L.) Sleumer, *Andira inermis* (W. Wright) H.B.K., and *Cecropia*



Figure 1—The trunk of a guamá, *Inga fagifolia* (L.) Willd., tree growing in Puerto Rico.

John K. Francis is a research forester at the International Institute of Tropical Forestry, U.S. Department of Agriculture, Forest Service, Río Piedras, PR 00928-2500; in cooperation with the University of Puerto Rico, Río Piedras, PR 00936-4984.




Figure 2.—The native range of *guamá*, *Inga fagifolia* (L.) Willd., in the West Indies is indicated by the shaded area.

schreberiana, are the principal species composing the secondary forests arising from abandoned shaded coffee plantations (28). Additional examples of species associated with *guamá* can be found in 2, 3, 6, 18, and 27.

LIFE HISTORY

Reproduction and Early Growth

Flowering and Fruiting.—The flower spikes are white and brushlike, 8 to 15 cm long, and borne singly or in pairs. The flat pods, 6 to 12 cm long and 2 to 3 cm wide, contain several seeds (16). *Guamá*, as a species, flowers and fruits throughout the year (16), although individual trees may be episodal in flowering and fruiting.

Seed Production and Dissemination.—Seeds for propagation may be collected by clipping ripe pods from trees or by picking them up from the ground under bearing trees. As pods ripen, they change color from green to yellow to brown. The testa of the seed is thin and leathery. The seeds will not retain their viability if dried. There were an average of 1,120 seeds per kg of fresh seeds in one sample from Puerto Rico (12). It is best to sow the seeds immediately after collection, but if this is not possible, the whole pods may be refrigerated for a few days without damage to the seeds. The seeds are dispersed by bats and birds that feed on the pods and/or the small amount of pulp surrounding the seeds. These animals may, in the process of feeding, carry the pods some distance from the mother tree (9, 20).

Seedling Development.—Germination is hypogeal. In one trial, germination began about 5 days after sowing and continued for 17 days (author, personal observation). Germination rates of 96 percent were observed in Puerto Rico (12), and rates of 80 percent, in Trinidad and Tobago (20). After germination under shade, the stems elongate about 6 cm before fully developing the first pair of leaves (author, personal observation). Grown under shade, a group of containerized seedlings reached an average height of 22 cm in 8 months after sowing. That nursery period could probably be

shortened by growing seedlings in full sun. Because about 6 months elapse before rapid growth begins in out-planted seedlings, it is advisable to use seedlings that are at least 50 cm in height to reduce the danger of the seedlings being smothered by weeds.

Vegetative Reproduction.—Small *guamá* trees will coppice, but large trees will not (20). It is not known whether cuttings can be rooted or if grafting can be successfully performed.

Sapling and Pole Stage to Maturity

Growth and Yield.—A solid planting of 52 potted seedlings in a small opening in moist forest grew an average of 1 m in the first 12 months, and another meter in the next 9 months. Survival was 98 percent at the end of 1 year and 96 percent at 21 months (author, personal observation). *Guamá* trees measured in late secondary forests in Puerto Rico showed mean annual diameter increments of about 0.5 cm per year over an 18-year period. Mean annual diameter increments for individual trees varied from undetectable to 1.2 cm (7). The largest *guamá* known in Puerto Rico measures 89.3 cm in d.b.h. and 21.5 m in height.¹ It is a former shade tree in an abandoned coffee plantation in the Luquillo Mountains, and it is at least 60 years old.

Rooting Habit.—Because of its deep and intertwining root habit, *guamá* was recommended for streambank stabilization in the Dominican Republic (22). *Guamá* roots are endomycorrhizal, and they produce nodules, presumably associated with *Rhizobium* bacteria (10).

Reaction to Competition.—*Guamá* is intermediate in tolerance to intolerant of shade. Seedlings will survive light to medium shade. Seedlings established by direct seeding under shade averaged 36 and 76 cm in height after 1 and 2 years, respectively. Corresponding heights of seedlings grown in full sun averaged 40 and 122 cm after 1 and 2 years (20). Adult trees retain vigor surprisingly well in lower canopy positions. *Guamá* trees in dominant, codominant, intermediate, and suppressed crown positions in late secondary, subtropical wet forest grew 0.77, 0.52, 0.33, and 0.19 cm/yr in d.b.h., respectively, over an 18-year period (7). In one experiment in Puerto Rico, two thinnings over an 18-year period resulted in an increase in average annual diameter increment from 0.38 to 0.47 cm/yr for *guamá* in secondary forest stands (26).

Although widely distributed, *guamá* normally constitutes only a small percentage of the total basal area of the forests where it grows. In a survey of the timber of privately owned secondary forests in moist portions of Puerto Rico, *guamá* ranked fifth of the species encountered in terms of basal area and accounted for 1.2 percent of the total basal area (4).

It is sometimes necessary to remove *guamá* trees because of poor form or low market value during timber stand improvement and cleaning operations. In a test of 46 trees, 80 percent were dead a year after frill girdling and treatment with 2,4,5-trichlorophenoxyacetic acid (2,4,5-T)² in diesel oil (23).

¹Champion tree register of Puerto Rico. On file with: International Institute of Tropical Forestry, U.S. Department of Agriculture, Forest Service, Río Piedras, PR 00928-2500.

²This herbicide is banned for silvicultural use in the United States by the U.S. Environmental Protection Agency.

Damaging Agents.—The most serious pest of guamá in Puerto Rico is the hormiguilla, an ant (*Myrmelachista ambigua ramulorum* Wheeler) that tunnels in live branches and trunks. This ant tends aphid species *Pseudococcus citri* Risso and *Cryptostigma inquilina* Newstead that suck the sap of the host tree and secrete a sweet nectar (19). Mortality was so high and growth so slow that in the 1920's and 1930's it caused a decline in coffee yields across Puerto Rico and resulted in growers switching to other species for coffee shade. There are great numbers of other insects that feed on guamá, generally with insignificant effects (21). A wilt disease called "mal de guaba," serious in *I. vera*, occasionally attacks guamá (19).

A survey of sawtimber on privately owned secondary forest lands in Puerto Rico found 38 percent of the guamá sawtimber free of defects, 35 percent with poor form, and 12 percent with stem cankers. It was estimated that only 8 percent of the wood volume was unsound (rotten) (1). Guamá logs are quickly attacked by pinhole borers, but resist sap-stain fungi (17). The wood in use is susceptible to termites and decay. The service life of untreated guamá posts is 1.5 to 1.6 years, but treated with pentachlorophenol³ in diesel oil, the service life is extended to 6 to 10 years (8, 11).

SPECIAL USES

At one time, guamá was the most popular shade tree for coffee production in Puerto Rico. Because of severe attacks by the hormiguilla, other shade tree species, particularly *I. vera*, are now favored (19). The small amount of white pulp surrounding the seeds is very sweet and is sometimes eaten by children. Livestock readily consume the pods (14). The flowers are an important source of nectar for honeybees (16).

One of the reasons that guamá was extensively planted as coffee shade is because it fixes nitrogen through symbiotic association with *Rhizobium* bacteria in its root nodules. It has also been shown that nitrogen fixation takes place through an unknown mechanism in the lenticular bark (29).

Guamá heartwood is pale reddish brown, often streaked with darker brown. The sapwood is not clearly distinguished from the heartwood. Samples of heartwood of six guamá trees in Puerto Rico averaged oven-dried weights of 0.64 ± 0.02 g/cm³ (author, personal observation). Guamá wood seasons rapidly with moderate degrade, shrinking 1.6 percent tangentially and 2.7 percent longitudinally (17). It works easily and finishes well. Guamá wood is suitable for furniture, cabinets, veneer, construction, and flooring, but because in Puerto Rico there are few logs of any species sawed, it is seldom used for any of these purposes. Guamá is used occasionally to make charcoal and posts.

GENETICS

Inga is a genus of about 200 species in tropical America (13). *Inga laurina* (Sw.) Willd. is a botanical synonym for guamá (15).

³The production and use of pentachlorophenol is now banned in the United States. Other treatments would probably result in somewhat shorter service lives.

LITERATURE CITED

1. Anderson, Robert L.; Birdsey, Richard A.; Barry, Patrick J. 1982. Incidence of damage and cull in Puerto Rico's timber resource, 1980. Resour. Bull. SO-88. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 13 p.
2. Beard, J.S. 1949. The natural vegetation of the Windward and Leeward Islands. Oxford, UK: Clarendon Press. 192 p.
3. Birdsey, Richard A.; Jiménez, Diego. 1985. The forests of Toro Negro. Res. Pap. SO-222. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 29 p.
4. Birdsey, Richard A.; Weaver, Peter L. 1982. The forest resources of Puerto Rico. Resour. Bull. SO-85. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 59 p.
5. Bisse, Johannes. 1981. Arboles de Cuba. Havana, Cuba: Editorial Científico-Técnica. 384 p.
6. Chinae, Jesús Danilo. 1980. The forest vegetation of the limestone hills of northern Puerto Rico. Ithaca, NY: Cornell University. 70 p. Ph.D. dissertation.
7. Crow, T.R.; Weaver, P.L. 1977. Tree growth in a moist tropical forest of Puerto Rico. Res. Pap. ITF-22. Río Piedras, PR: U.S. Department of Agriculture, Forest Service, Institute of Tropical Forestry. 17 p.
8. Chudnoff, M.; Maldonado, E.D. 1964. Preservative treatments and service life of fence posts in Puerto Rico. Res. Pap. ITF-1. Río Piedras, PR: U.S. Department of Agriculture, Forest Service, Institute of Tropical Forestry. 31 p.
9. Devoe, Nora Nancy. 1989. Differential seeding and regeneration in openings and beneath closed canopy in subtropical wet forest. New Haven, CT: Yale University. 307 p. Ph.D. dissertation.
10. Edmisten, Joe. 1970. Survey of mycorrhiza and nodules in the El Verde Forest. In: Odum, Howard T.; Pigeon, Robert. A tropical rain forest. Washington, DC: U.S. Atomic Energy Commission, Division of Technical Information: F15-F20.
11. Englerth, George H. 1960. The service life of untreated posts in Puerto Rico after one year in test. Trop. For. Notes 5. Río Piedras, PR: U.S. Department of Agriculture, Forest Service, Tropical Forest Research Center. 2 p.
12. Francis, John K.; Rodríguez, Alberto. 1993. Seeds of Puerto Rican trees and shrubs: second installment. Res. Note SO-374. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 5 p.
13. Howard, Richard A. 1988. Flora of the Lesser Antilles: Leeward and Windward Islands. Jamaica Plain, MA: Harvard University, Arnold Arboretum. 673 p. Vol. 4.
14. Liogier, Henri Alain. 1978. Arboles Dominicanos. Santo Domingo, Dominican Republic: Academia de Ciencias de la Republica Dominicana. 220 p.
15. Liogier, Henri Alain; Martorell, Luís F. 1982. Flora of Puerto Rico and adjacent islands: a systematic synopsis. Río Piedras, PR: Editorial de la Universidad de Puerto Rico. 342 p.

16. Little, Elbert L., Jr.; Wadsworth, Frank H. 1964. Common trees of Puerto Rico and the Virgin Islands. Agric. Handb. 249. Washington, DC: U.S. Department of Agriculture. 548 p.
17. Longwood, Franklin R. 1961. Puerto Rican woods, their machining, seasoning and related characteristics. Agric. Handb. 205. Washington, DC: U.S. Department of Agriculture. 98 p.
18. Mackler, R.; Hannah, P. 1988. The forest at Wallings Reservoir, Antigua, West Indies, 44 years after J.S. Beard's study: conjecture on the patterns of change. Commonwealth Forestry Review. 67(3): 269-278.
19. Marrero, José. 1954. Especies del genero *Inga* usadas como sombra de cafe en Puerto Rico. Caribbean Forester. 15(1/2): 54-71.
20. Marshall, R.C. 1939. Silviculture of the trees of Trinidad and Tobago, British West Indies. London: Oxford University Press. 247 p.
21. Martorell, Luis F. 1975. Annotated food plant catalog of the insects of Puerto Rico. Río Piedras, PR: University of Puerto Rico, Agricultural Experiment Station. 303 p.
22. Schiffino, José. 1945. Riqueza forestal Dominicana. Trujillo, Dominican Republic: Editorial Montalvo. 291 p. Vol. 1.
23. Sposta, Joseph W. 1960. Chemical removal of inferior tropical tree species. Trop. For. Notes 4. Río Piedras, PR: U.S. Department of Agriculture, Forest Service, Tropical Forest Research Center. 2 p.
24. Veillon, Jean Pierre. 1986. Especies forestales autóctonas de los bosques naturales de Venezuela. Mérida, Venezuela: Instituto Forestal Latinoamericano, Universidad de los Andes. 199 p.
25. Wadsworth, Frank H. 1951. Forest management in the Luquillo Mountains. 1: The setting. Caribbean Forester. 12(3): 93-114.
26. Weaver, Peter L. 1983. Tree growth and stand changes in the subtropical life zones of the Luquillo Mountains of Puerto Rico. Res. Pap. SO-190. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 24 p.
27. Weaver, Peter L. 1991. Environmental gradients affect forest composition in the Luquillo Mountains of Puerto Rico. Interciencia. 16(3): 142-151.
28. Weaver, P.L.; Birdsey, R.A. 1986. Tree succession and management opportunities in coffee shade stands. Turrialba. 36(1): 47-58.
29. Yatazawa, M.; Hambali, G.G.; Uchino, F. 1983. Nitrogen fixing activity in warty lenticilate tree barks. Soil Science and Plant Nutrition. 29(3): 285-294.

UNIVERSITY OF MINNESOTA



3 1951 D02 980 825 0

This publication reports research involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.